

Using Naloxone

- Reverses opioid overdose
- Safe to use
- Works in 2–5 minutes
- May need more than one dose



How to respond to an overdose




- 1. Call 911**
Tell them someone is not breathing.

- 2. Give naloxone**
Do not test device.
Each device can only be used once.

Peel



Place
Lay person on their back and tilt their head up.



Press
Spray in one nostril by pushing plunger.





- 3. Recovery position**
Place the person on their side to prevent choking.

- 4. Repeat**
Give second dose if no response in two minutes.



- 5. Stay with the person**
Wait until medical help arrives.

Uso de Naloxona

- Revierte la sobredosis de opioides
- Seguro de usar
- Funciona en 2 a 5 minutos
- Puede ser necesaria más de una dosis



Cómo responder ante una sobredosis




1. Llame al 911
Dígales que alguien no está respirando.

2. Administre la naloxona
No pruebe el dispositivo.
Cada dispositivo solo puede usarse una vez.


Abra



Coloque
Acueste a la persona boca arriba e incline la cabeza hacia arriba.



Presione
Rocíe en una fosa nasal presionando el émbolo.




3. Posición de recuperación
Coloque a la persona de lado para evitar que se asfixie.

4. Repita
Administre una segunda dosis si no respuesta en dos minutos.



5. Permanezca con la persona
Espere hasta que llegue la ayuda médica.




Call 911

Tell them someone is not breathing.


Peel

Do not test device. Each device can only be used once.



Place

Lay person on their back and tilt their head up.



Press

Spray in one nostril by pushing plunger.

